



The NERD Dataset: Near East Radiocarbon Dates between 15,000 and 1,500 cal. yr. BP

ALESSIO PALMISANO

ANDREW BEVAN

DAN LAWRENCE

STEPHEN SHENNAN

**Author affiliations can be found in the back matter of this article*

DATA PAPER

][ubiquity press

ABSTRACT

To our knowledge, the dataset described in this paper represents the largest existing repository of uncalibrated radiocarbon dates for the whole Near East from the Late Pleistocene to the Late Holocene (15,000 – 1,500 cal. yr. BP). It is composed of 11,027 radiocarbon dates from 1,023 sites that have been collected comprehensively by cross-checking multiple sources (extant digital archives and databases, edited volumes, monographs, journals papers, archaeological excavation reports, etc.) under the umbrella of the Leverhulme Trust funded project “Changing the Face of the Mediterranean” and of the ERC project “CLASS – Climate, Landscape, Settlement and Society: Exploring Human-Environment Interaction in the Ancient Near East”. This is an ongoing dataset that will be updated step by step with newly published radiocarbon dates.

CORRESPONDING AUTHORS:

Alessio Palmisano

Department of Ancient History,
Ludwig Maximilian University
of Munich, Munich, DE

alessio.palmisano@lrz.uni-muenchen.de

Andrew Bevan

Institute of Archaeology,
University College London,
London, GB

a.bevan@ucl.ac.uk

KEYWORDS:

Radiocarbon dates; Near East;
Holocene; Archaeological sites;
Late Pleistocene; Mesopotamia

TO CITE THIS ARTICLE:

Palmisano A, Bevan A,
Lawrence D, Shennan S. 2022
The NERD Dataset: Near East
Radiocarbon Dates between
15,000 and 1,500 cal. yr. BP.
*Journal of Open Archaeology
Data*, 10: 2, pp. 1–9. DOI:
<https://doi.org/10.5334/joad.90>

(1) OVERVIEW

CONTEXT

The present dataset was created under the scope of the Leverhulme Trust funded project “Changing the Face of the Mediterranean” (grant number: RPG-2015-031, PI Neil Roberts) as a part of two regional case studies ([1, 2], for the special issue see [3]) and of the ERC project “CLASS – Climate, Landscape, Settlement and Society: Exploring Human-Environment Interaction in the Ancient Near East” (grant number: 802424, PI Dan Lawrence). Both projects explore trends in human-environment interactions respectively in the Mediterranean basin and the Near East between Late Pleistocene and Late Holocene. In these two projects, long-term demographic trends have been compared with climatic shifts and vegetation change in order to highlight possible causal relationships between them.

In this context, large lists of radiocarbon dates from archaeological contexts can be calibrated and counted up (summed in the manner of a histogram) as a proxy for population, based on the assumption that the more people living in a given region, the more the archaeological deposits, the more organic materials, and the more radiocarbon samples collected and dated. As a consequence, summed probability distributions (SPDs) of calibrated radiocarbon dates have become particularly popular for inferring demographic trends in prehistory and for assessing population response to climate shifts and the anthropic impact on landscapes [4–10]. The radiocarbon dates provide a better chronological resolution than archaeo-demographic proxies based

on typo-chronological schemes (e.g., raw site count, aggregated estimated settlement size, number of burials, count of potsherds), where reliance on relatively long-lived artefactual data types means phases may span several centuries. Hence, comparing radiocarbon-inferred demographic trends with palaeoclimatic records and fossil pollen cores offers further opportunities to understand long-term socio-ecological trajectories.

Although radiocarbon dates from the Near East are available in some extant digital repositories and databases, they face several challenges, such as inconsistent chronological and geographical coverage, missing or inaccurate spatial location, missing updates from recent publications, lack of interoperable file formats, and unspecified licenses. These issues hamper the interoperability and the reuse of radiocarbon dates for comparative studies and do not adhere to the FAIR (findable, accessible, interoperable, reusable) principles for scientific data management and stewardship [11]. Comparisons of long-term demographic trends across the whole Near East require a synthetic dataset of radiocarbon dates collated under the same protocol easily accessible to a wider audience.

Here we provide the largest list so far of published radiocarbon dates for the whole Near East from the Late Pleistocene to Late Holocene between 15,000 and 1,500 cal. yr. BP. A total of 11,027 sites from 1,023 archaeological sites have been collated and harmonised from extant online digital repositories, original digital and print publications and archaeological excavation reports (*Figure 1*). The NERD dataset is hosted in the collaborative



Figure 1 Map of the Near East showing the spatial distribution of radiocarbon samples.

online platform GitHub, where registered users can contribute to the content of the original repository. Therefore, NERD offers a novel dynamic resource that can be updated at any time with new radiocarbon dates without the restrictions and limitations of traditional static databases.

Spatial coverage

Description: The dataset spatially covers the whole Near East which is approximately 5,900,000 sq. km ([Figure 1](#)). The radiocarbon dates are distributed over the following countries: Armenia, Azerbaijan, Bahrain, Cyprus, Egypt, Georgia, Israel, Iraq, Jordan, Kuwait, Palestine, Russia, Saudi Arabia, Syria, Turkey, United Arab Emirates, and Yemen.

The decimal degree coordinates of the minimum-bounding box are given in the geographic coordinate system WGS 84.

Northern boundary: + 42.7400
Southern boundary: + 13.0204
Eastern boundary: + 61.3377
Western boundary: + 25.7084

Temporal coverage

Dates range from the Late Pleistocene to the Late Holocene (15,000 – 1,500 cal. yr. BP/13,000 BC – 500 AD).

(2) METHODS

The creation of this dataset was eased by the availability of openly accessible digital online repositories and by a growing consensus among scholars about the benefits of data sharing to advance knowledge [12]. The collation of all radiocarbon dates included in the present dataset was enabled by a desk-based synthesis integrating and harmonising secondary published sources and extant online databases.

STEPS

The radiocarbon dates were collected in four different steps. Firstly, we extracted and merged uncalibrated radiocarbon dates from existing online digital archives and databases (ArAGATS project: [13]; BANADORA: [14]; [15]; CalPal: [16]; CONTEXT: [17]; [18]; IRPA/KIK: [19]; ORAU: [20]; PPNL: [21]; RADON: [22]; TAY project: [23]; 14SEA: [24]).

Secondly, we enhanced the list of radiocarbon dates by a comprehensive screening of all the most relevant journals available online up to November 2021. Further dates were extracted from several monographs, chapters in edited volumes, archaeological excavation reports, and websites. We have not been able to include unpublished dates or those published in non-digital grey literature. The total number of references scrutinised to

build our dataset is 719. As a result, we collected 11,027 radiocarbon dates from 1,023 archaeological sites. To our knowledge, this is the largest collation of published radiocarbon dates for the whole Near East.

Thirdly, we standardised the data and comparatively cross-checked all spatial coordinates from extant online digital archives and publications to guarantee maximum accuracy in the geographic location.

Lastly, a unique SiteID has been assigned to each radiocarbon date in order to link all of them to a specific site of provenance. Wherever possible, all information related to the radiocarbon samples has been recorded such as “LabID”, “Material”, and “Species”.

SAMPLING STRATEGY

The dataset derives from existing publications relevant to the chronological scope of interest, spanning from the Late Pleistocene to the Late Holocene (15,000 – 1,500 cal. yr. BP). We tried to record all the known radiocarbon dates published until November 2021 that fall within our chronological scope of interest.

QUALITY CONTROL

After the data collection stage, we cleaned and checked all the information inputted, as the collated radiocarbon dates come from various and inconsistent secondary sources. All radiocarbon dates have been checked for duplicates and have been standardised. For instance, some radiocarbon dates come from sites that have multiple site names and we assigned them a unique descriptive field “SiteID” and standardised the “SiteName” to avoid duplication of sites. Likewise, we have checked duplicated radiocarbon laboratory codes and removed those that refer to the same raw date. The “LabID” was standardised according to the following list: https://radiocarbon.webhost.uits.arizona.edu/sites/default/files/Labs-2021_09_03.pdf. For those radiocarbon dates having more than one lab ID, we created a second field named “OthLabID”, which provides the alternative lab ID.

In the field “Source”, we reported the original publication of each date and all subsequent works. During the data cleaning stage, we looked for any possible inconsistencies in the publications that could result in conflicting statements in the attributes (e.g., conventional pre-calibration radiocarbon age (“CRA”), 1-standard deviation error (“Error”), the material of the organic sample (“Material”), duplicated lab ID) associated with the radiocarbon dates. When we encountered discrepancies in the sources, the final entry in our dataset was based on the most detailed descriptions, on the original publication, on a comparison between multiple later publications and on any more recent measurement revision. Any issue associated with a specific date is reported in the field “Problems” and a more extensive description is recorded in the “Comments” field.

The geographic locations of the radiocarbon samples (“Longitude” and “Latitude”) were assigned in decimal degrees according to the World Geodetic System 1984 (WGS84, EPSG:4326). The location quality (“LocQual”) has been ranked according to five levels of accuracy: A (centroid of the archaeological site from which the organic sample has been collected), B (within +/- 2 km), C (within +/- 5 km), D (within +/- 10 km), E (within +/- 20 km). Most dates have the highest level of accuracy ($n = 8851$, category A), while only 129 have a coarse level of spatial accuracy in categories D and E. The radiocarbon dates have been systematically georeferenced by making use of any possible sources such as extant digital repositories, archaeological survey maps, satellite images, websites, coordinates from original publications, and digital tools (e.g., Google Earth). The coordinates from extant online digital archives (e.g., RADON, CalPal; BANADORA, etc.) have been manually cross-checked to avoid any possible spatial discrepancy and ambiguity due to the fact that it is unclear if those coordinates represent the centroid of the archaeological sites, or of the closest towns or villages. Radiocarbon samples with no geographic coordinates or unclear spatial location have not been included in the dataset. We are aware that publishing the geographic coordinates could make archaeological sites more vulnerable to looting activities. However, we did not obfuscate the locations of radiocarbon dates because they come from excavated archaeological sites whose coordinates are widely available through unrestricted resources publicly accessible (e.g., published archaeological surveys reports, monographies, digital publications, geographic tags in Google Earth, existing online databases).

CONSTRAINTS

Although we have undertaken a scrupulous effort in data cleaning and checking, the present dataset may still contain some errors possibly inherited from multiple resources that were screened or due to human error. Hence, we would encourage all users of the NERD dataset to inform us of any errors so that we can correct them and update the dynamic online repository.

There are several caveats when dealing with radiocarbon dates. Unlike other areas of the world where archaeological work is routinely integrated into planning and construction industries through a long tradition of regulated commercial archaeology (e.g., North America, central and northern Europe), in southwestern Asia most archaeological investigations are carried out by academic projects. As a consequence, this region is more prone to spatial and chronological biases due to investigators interests (e.g., the spread of farming, the human response to rapid climatic shifts such as the 4.2 and 3.2 kya events) than areas with a long tradition of commercial archaeology. Regarding chronological biases, radiocarbon dates are particularly prevalent in the Late

Pleistocene and Early Holocene due to the absence of high-resolution typo-chronological schemes. Conversely, for later historical periods archaeologists rely more on short-lived pottery types and historical media (e.g., clay tablets, seals and coins) for dating archaeological layers rather than using radiometric dating. A certain reluctance in using radiocarbon dating among those archaeologists digging Iron Age sites is also justified by the Hallstatt radiocarbon calibration plateau (ca. 2750–2350 cal. yr. BP) which makes it difficult to obtain refined radiocarbon-based chronologies. On a pan-regional scale, the present dataset guarantees a good chronological coverage until ~3000 cal. yr. BP, while other regions such as Mesopotamia, Anatolia and Iran show a research-biased drop in the available radiocarbon dates from 4000–3500 cal. yr. BP onwards as shown recently by Palmisano et al [9].

Regarding spatial bias, 35% of the radiocarbon dates come from the Levant (modern Israel, Palestine, Lebanon, Jordan and western Syria), while other regions such as Iran, Arabia and southern Mesopotamia are underrepresented (*Figure 2*). Furthermore, 70% of the Levantine dates come from modern Israel and Palestine. This is due to the higher research budgets of Israeli archaeological teams interested in producing or improving an absolute chronology.

Additionally, certain chronological periods are more likely to be sampled than others. A clear research bias in the southern Levant is due to the interest of many archaeologists in providing a better chronology for the Early Bronze Age sub-periods (ca. 3800–2500 BC) and the Late Bronze Age/Iron Age transition (ca. 1200–950 BC) [25, 26].

The radiocarbon dates have been collected from a variety of existing sources with varying levels of published information. For instance, not all samples have published information about the site context, sample material, sample species, etc., although we tried to recover as much missing information as possible. For example, 87% of the listed dates have information about the sample material (e.g., charcoal, bone, wood, etc.), but only 28% have information about species (e.g., sample of wheat seeds, olive stones, etc.). Most of our dataset is composed of long-lived unidentified charcoal samples that could be affected by the so-called old wood effect. Their use could affect the resulting summed probability distributions (SPDs) of calibrated radiocarbon dates that have been widely used by palaeo-demographers for modelling past population human dynamics. To explore this issue, we produced SPDs of unnormalised calibrated radiocarbon dates from short-lived radiocarbon samples (e.g., bones, collagen, seeds, grains) and from all dates. The SPDs including all dates (*Figure 3*, in grey) and only short-lived dates (*Figure 3*, in green) are highly correlated ($r = 0.93$, p -value = 0.01, Pearson) and the so-called old wood effect seems not to have significantly affected the resulting post-calibration probability densities.




Figure 2 Kernel density map of radiocarbon samples.




Figure 3 Summed probability distributions of unnormalised calibrated radiocarbon dates including all samples (in grey) and the short-lived samples (in green).

(3) DATASET DESCRIPTION

The dataset contains a set of three files providing a list of radiocarbon dates (*nerd.csv*), a file providing a list of references (*References.txt* file) of published radiocarbon dates stored in the *nerd.csv* file (see field “Source”), and a metadata and field description (see [Table 1](#)) for the attributes of the radiocarbon samples (*README.md*).

The NERD dataset has been published in accordance with the FAIR principles. There are several steps that need to be implemented before making our data easily findable and reusable (see [Figure 4](#)). After storing all radiocarbon dates into a *.csv* file, we published it publicly via GitHub, which is an ideal platform to modify, delete, and rewrite any content of a project but not for lasting referencing purposes. For this latter purpose, Zenodo

DATAFIELD	DESCRIPTION
DateID (numeric)	unique identifier for the radiocarbon sample
LabID (character)	unique identifier for the lab's radiocarbon sample
OthLabID (character)	unique alternative identifier for the lab's radiocarbon sample
Problems (character)	problems related to the radiocarbon sample (e.g. missing Lab Id, duplicated Lab Ids, etc.)
CRA (numeric)	radiocarbon concentration expressed in years before present (BP)
Error (numeric)	Standard error of radiocarbon date in years
DC13 (numeric)	Isotopic fractionation of stable carbon isotopes Carbon-13
Material (character)	material of the radiocarbon sample
Species (character)	species of the radiocarbon sample
SiteID (numeric)	unique identifier of the site from which the radiocarbon sample has been collected
SiteName (character)	name of archaeological site
SiteContext (character)	original archaeological context from which the radiocarbon sample was collected
SiteType (character)	type of archaeological site
Country (character)	country from which the radiocarbon sample was collected
Longitude (numeric)	WGS84 eastings
Latitude (numeric)	WGS84 northings
LocQual (character)	scale defining the accuracy of the spatial coordinates of radiocarbon samples
Source (character)	source from which the radiocarbon samples have been collected
Comment (character)	Comments about the issues reported in the field "Problems"

Table 1 Description of the fields in the file nerd.csv.




Figure 4 The road map to the FAIR principles (findable, accessible, interoperable, reusable).

offers more security for permanently archiving a specific dataset by assigning a Digital Object Identifier (DOI), which makes the repository more findable and citable. Hence, we authorized the GitHub account of the leading author to be connected to Zenodo. This allows Zenodo to publish online the repository stored on Github by issuing a DOI. This step is possible by creating a release in GitHub. Each time we created a new version of the repository,

we created a new release that can be tracked in the repository on Github. Thanks to this step, any version of the repository is stored both on Github and Zenodo. The strength of this approach is to make our dataset dynamic and easy to be permanently stored online. In addition, the collaborative platform Github allows any registered user to contribute to NERD by adding new radiocarbon dates not included in our dataset through the command

‘Pull request’. The final stage of the roadmap to the FAIR principles is to publish a data paper of NERD in order to make it more visible to the research community.

Finally, NERD is also currently integrated into the R package *c14bazAAR*, which allows for the querying, managing and merging of various open-access radiocarbon archives [27]. This makes NERD more accessible and usable within the scientific community.

OBJECT NAME

nerd.csv
References.txt
README.md

DATA TYPE

Secondary data, and processed data from originally published materials.

FORMAT NAMES AND VERSIONS

.csv, .txt, .md

CREATION DATES

The dataset was created in three different stages. 70% of the dataset was created during 2016–2018 as part of the Leverhulme Trust funded project “Changing the face of the Mediterranean: land cover and population since the advent of farming”. The remaining portion of the dataset was created between October 2019 and December 2021 under the umbrella of the ERC project “CLASS – Climate, Landscape, Settlement and Society: Exploring Human-Environment Interaction in the Ancient Near East” and of an Alexander von Humboldt Research Fellowship.

DATASET CREATORS

The researcher responsible for the data entry and management was Alessio Palmisano. A portion of radiocarbon dates for the Levant were added by Fayrouz Ibrahim and Alexander Kabelindde. Andrew Bevan was also heavily involved in checking and restructuring parts of the dataset.

LANGUAGE

English

LICENSE

Creative Commons License CC-BY 4.0: <https://creativecommons.org/licenses/by/4.0/>

REPOSITORY LOCATION

<https://doi.org/10.5281/zenodo.5767862>

GITHUB LOCATION

<https://github.com/apalmisano82/NERD>

PUBLICATION DATE

08/12/2021

(4) REUSE POTENTIAL

This data set represents what is to our knowledge the largest existing collation of radiocarbon dates for the whole Near East from the Late Pleistocene to the Late Holocene (15,000 – 1,500 cal. yr. BP).

The locations of the archaeological sites yielding radiocarbon samples have been collected guaranteeing the highest possible accuracy and represent a basic resource for georeferencing of past Near Eastern settlements. This large dataset can be added with other lists of radiocarbon dates for comparative studies and research agendas focussing on a broader geographical scope. The anthropogenic radiocarbon dates can be used as a proxy of human population for building summed probability distributions (SPDs), an established technique for inferring past population fluctuations across space and time. The large size of the present dataset and its structure make it suitable for a wide range of spatio-temporal analyses, with the potential for careful handling of the underlying temporal uncertainty [28, 29]. The present dataset can be integrated with archaeological artefacts, and architectural remains for revising established typo-chronological schemes and building new ones. Furthermore, the combination of this large list of radiocarbon dates with other scientific data, such as fossil pollen cores, paleoclimatic records, and archaeological survey data, opens new comparative perspectives in the study of long-term human-environment interactions at different chronological and spatial scales of analyses.

ACKNOWLEDGEMENTS

We thank Fayrouz Ibrahim and Alexander Kabelindde for helping with the inputting and georeferencing of the Levantine radiocarbon dates. We also thank the three anonymous reviewers for their comments on the earlier version of the paper.

FUNDING INFORMATION

The archive NERD has been collated thanks to the following grants: 1) Leverhulme Trust project “Changing the Face of The Mediterranean: Land cover and population since the advent of farming” (grant number RPG-2015-031, award holders: Neil Roberts (PI), Andrew Bevan (Co-I), Ralph Fyfe (Co-I), & Stephen Shennan (Co-I)); 2) the European Research Council under the European Union’s Horizon 2020 research and innovation programme project “CLaSS – Climate, Landscape, Settlement and Society: Exploring Human Environment Interaction in the Ancient Near East” (grant number 802424, award holder: Dan Lawrence); 3) Alexander von Humboldt Research Fellowship (award holder: Alessio Palmisano).

COMPETING INTERESTS

The leading author is a co-chief editor for *Journal of Open Archaeology Data*, which is on a voluntary basis. However, he did not participate in any part of the editorial process and another member of the team have had assigned responsibility for overseeing peer review. All authors of the paper declare that they have no competing interests.

AUTHOR AFFILIATIONS

Alessio Palmisano  orcid.org/0000-0003-0758-5032

Department of Ancient History, Ludwig Maximilian University of Munich, Munich, DE

Andrew Bevan  orcid.org/0000-0001-7967-3117

Institute of Archaeology, University College London, London, GB

Dan Lawrence  orcid.org/0000-0001-5613-1243

Department of Archaeology, Durham University, Durham, GB

Stephen Shennan  orcid.org/0000-0001-6605-064X

Institute of Archaeology, University College London, London, GB

REFERENCES

- Palmisano A, Woodbridge J, Roberts N, Bevan A, Fyfe R, Shennan S, Cheddadi R, Greenberg R, Kaniewski D, Langgut D, Leroy S, Litt T, Miebach A.** Holocene Landscape Dynamics and Long-term Population Trends in the Levant. *The Holocene*, 2019; 29(5): 708–727. DOI: <https://doi.org/10.1177/0959683619826642>
- Woodbridge J, Roberts N, Palmisano A, Bevan A, Shennan S, Fyfe R, Eastwood WJ, Izdebski A, Cakirlar C, Woldring H, Broothaerts N, Kaniewski D, Finn M, Labuhn I.** Pollen-inferred regional vegetation patterns and demographic change in Southern Anatolia through the Holocene. *The Holocene*, 2019; 29(5): 728–741. DOI: <https://doi.org/10.1177/0959683619826635>
- Bevan A, Palmisano A, Woodbridge J, Fyfe R, Shennan S, Roberts N.** The changing face of the Mediterranean: land cover, demography and environmental change: Introduction and overview. *The Holocene*, 2019; 29(5): 703–707. DOI: <https://doi.org/10.1177/0959683619826688>
- Rick JW.** Dates as data: an examination of the Peruvian preceramic radiocarbon record. *American Antiquity*, 1987; 52(1): 55–73. DOI: <https://doi.org/10.2307/281060>
- Shennan S, Downey S, Timpson A,** et al. Regional population collapse followed initial agriculture booms in mid-Holocene Europe. *Nature Communications*, 2013; 4: 2486. DOI: <https://doi.org/10.1038/ncomms3486>
- Bevan A, Colledge S, Fuller D, Fyfe R, Shennan S, Stevens C.** Holocene fluctuations in human population demonstrate repeated links to food production and climate. *Proceedings of the National Academy of Sciences*, 2017; 114(49): E10524–E10531. DOI: <https://doi.org/10.1073/pnas.1709190114>
- Roberts N, Woodbridge J, Palmisano A, Bevan A, Fyfe R, Shennan S.** Mediterranean landscape change during the Holocene: synthesis, comparison and regional trends in population, land cover and climate. *The Holocene*, 2019; 29(5): 923–937. DOI: <https://doi.org/10.1177/0959683619826697>
- Crema ER, Bevan A.** Inference from large sets of radiocarbon dates: software and methods. *Radiocarbon*, 2021; 63(1): 23–39. DOI: <https://doi.org/10.1017/RDC.2020.95>
- Palmisano A, Lawrence D, de Gruchy MW, Bevan A, Shennan S.** Holocene regional population dynamics and climatic trends in the Near East: a first comparison using archaeo-demographic proxies. *Quaternary Science Reviews*, 2021; 252: 106739. DOI: <https://doi.org/10.1016/j.quascirev.2020.106739>
- Bird D, Miranda L, Vander Linden M, Robinson E, Bocinsky RK, Nicholson C, Capriles JM, Finley JB, Gayo EM, Gil A, d'Alpoim Guedes J, Hoggarth JA, Kay A, Loftus E, Lombardo U, Mackie M, Palmisano A, Solheim S, Kelly RL, Freeman J.** p3k14c, a synthetic global database of archaeological radiocarbon dates. *Scientific Data*, 2022; 9: 27. <https://doi.org/10.1038/s41597-022-01118-7>
- Wilkinson M, Dumontier M, Aalbersberg I,** et al. The FAIR Guiding Principles for scientific data management and stewardship. *Scientific Data*, 2016; 3: 160018. DOI: <https://doi.org/10.1038/sdata.2016.18>
- Bevan A.** The data deluge. *Antiquity*, 2015; 89(348): 1473–1484. DOI: <https://doi.org/10.15184/aqy.2015.102>
- ArAGATS Data Portal: <https://aragats.gorgesapps.us/>, accessed 10th November 2019.
- BANADORA. Banque Nationale de Données Radiocarbone pour l'Europe et le Proche Orient, Centre de Datation par le Radiocarbone [Internet]. Lyon: CNRS. Available from: <http://www.arar.mom.fr/banadora/>, accessed 31st May 2018.
- Borrell F, Junno A, Barceló JA.** Synchronous environmental and cultural change in the emergence of agricultural economies 10,000 years ago in the Levant. *PLoS one*, 2015; 10(8): e0134810. DOI: <https://doi.org/10.1371/journal.pone.0134810>
- Weninger B.** 2018. CalPal Archaeological 14C-Database. Available at: https://www.academia.edu/36766067/CalPal_Archaeological_14C-Database_Europe.xlsx_June_2018, Accessed 20th November 2019.
- Böhner U, Schyle D.** Radiocarbon CONTEXT database 2002–2006; 2006. Available at <http://context-database.uni-koeln.de/>.
- Flohr P, Fleitmann D, Matthews R, Matthews W, Black S.** Evidence of resilience to past climate change in Southwest Asia: Early farming communities and the 9.2 and 8.2 ka events. *Quaternary Science Reviews*, 2016; 136: 23–39. DOI: <https://doi.org/10.1016/j.quascirev.2015.06.022>
- Van Strydonck M, De Roock E.** Royal Institute for Cultural Heritage web-based radiocarbon database. *Radiocarbon*, 2011; 53(2): 367–370. DOI: <https://doi.org/10.1017/S0033822200056629>

20. **ORAU.** Oxford Radiocarbon Accelerator Unit online database; 2016. Oxford: University of Oxford. Available at: <https://c14.arch.ox.ac.uk/database/db.php>.
21. **Benz M.** 2014. PPND – the Platform for Neolithic Radiocarbon Dates. Available at: https://www.exoriente.org/associated_projects/ppnd.php.
22. **Hinz M, Furholt M, Müller J, Raetzel-Fabian D, Rinne C, Sjögren KG, Wotzka HP.** RADON - Radiocarbon dates online 2012. Central European database of ¹⁴C dates for the Neolithic and Early Bronze Age. *Journal of Neolithic Archaeology*, 2012; 14: 1–4. Available at: <http://radon.ufg.uni-kiel.de/>.
23. TAY project ¹⁴C Database (22nd Update), 2019. Available at: <http://tayproject.org/C14searcheng.html>.
24. **Reingruber A, Thissen L.** The 14SEA Project. A ¹⁴C database for Southeast Europe and Anatolia (10,000–3000calBC); 2017. Available at: http://www.14sea.org/2_dates.html.
25. **Finkelstein I, Piasetzky E.** The Iron Age Chronology Debate: Is the Gap Narrowing? *Near Eastern Archaeology*, 2011; 74: 50–54. DOI: <https://doi.org/10.5615/neareastarch.74.1.0050>
26. **Regev J, De Miroschedji P, Greenberg R, Braun E, Greenhut Z, Boaretto E.** Chronology of the Early Bronze Age in the southern Levant: new analysis for a high chronology. *Radiocarbon*, 2012; 54(3–4): 525–566. DOI: <https://doi.org/10.1017/S003382220004724X>
27. **Schmid C, Seidensticker D, Hinz M.** 2019. c14bazAAR: An R package for downloading and preparing C14 dates from different source databases. *Journal of Open Source Software*, 2019; 4(43): 1914. DOI: <https://doi.org/10.21105/joss.01914>
28. **Crema ER, Bevan A, Shennan S.** Spatiotemporal approaches to archaeological radiocarbon dates. *Journal of Archaeological Science*, 2017; 87: 1–9. DOI: <https://doi.org/10.1016/j.jas.2017.09.007>
29. **Bevan A, Crema E, Li X, Palmisano A.** Intensities, interactions and uncertainties: some new approaches to archaeological distributions. In: Bevan A, Lake M (eds.), *Computational Approaches to Archaeological Space Walnut*. 2013; 27–52. Creek: Left Coast Press.

TO CITE THIS ARTICLE:

Palmisano A, Bevan A, Lawrence D, Shennan S. 2022 The NERD Dataset: Near East Radiocarbon Dates between 15,000 and 1,500 cal. yr. BP. *Journal of Open Archaeology Data*, 10: 2, pp. 1–9. DOI: <https://doi.org/10.5334/joad.90>

Published: 22 February 2022

COPYRIGHT:

© 2022 The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC-BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. See <http://creativecommons.org/licenses/by/4.0/>.

Journal of Open Archaeology Data is a peer-reviewed open access journal published by Ubiquity Press.

